CUMULATIVE AUTHOR INDEX
Annals of Entomology
Volume 99, 2006

A

Abrahamson, W. G., 656
Akimoto, S.-I., 457
Alarie, Y., 743
Albuquerque, G. S., 7
Aldrich, J. R., 91
Almeida, R. P. P., 577, 884
Aluja, M., 662
Amir-Maafi, M., 84
Amrine, J. W., Jr., 1057
Anderson, K. G., 305
Anderson, R. S., 1258, 1261
Archibald, S. B., 487
Ark, J. T., 560
Armbruster, P., 1234
Ashe, J. S., 19
Austin, J. W., 697

B

Barbour, J. D., 718
Barclay, H. J., 607
Beal, A. L. H., 545
Benoit, J. B., 560
Bentz, B. J., 189
Berlocher, S., 662
Birge, L. M., 449
Blossey, B., 683
Boivin, G., 1121
Bomar, C. R., 1257
Bostanian, N. J., 536
Boyce, S. L., 164
Boyd, E. A., 1154
Branch, C. L., 1146
Branham, M. A., 409
Brannon, S. L., 300
Braswell, W. E., 449
Brelsfoard, C. L., 707
Broce, A. B., 157
Brower, A. V. Z., 184
Brunner, J. F., 169, 617
Bugrov, A. G., 457
Burand, J. P., 969
Buschman, L. L., 279
Butler, C. D., 317
Byers, J. A., 865

C

Caccone, A., 195
Cáceres, C., 387
Caceres, C., 1191
Cagnotti, C. L., 70
Carmichael, A., 567
Carroll, M. K., 697
Castro, B. A., 999
Catarino, M., 1260
Cattelan, A. J., 91
Causton, C. E., 121
Cave, R. D., 776, 1146
Cavichioli, R. R., 648
Cayol, J.-P., 387
Cervantes, D. E., 718
Challet, G. L., 743
Chang, C. L., 1191
Chase, J. A., 463
Chatzimanolis, S., 19
Chen, T.-Y., 917
Cheng, J.-A., 1127
Chi, H., 84
Cho, I. K., 1191
Choi, M.-Y., 905
Christiansen, P., 342
Chu, C.-C., 917
Chuang, Y.-C., 859
Cisneros, R., 945
Cladera, J. L., 70
Coelho, J. R., 793
Cognato, A. I., 1034
Conn, J. E., 1234
Conte, H., 1206
Cook, W. M., 809
Copeland, R. S., 261, 530
Costa, C., 409
Cover, S. P., 487
Crampton, L., 33
Creel, O. R., 1146
Crespi, B. J., 407
Cruz, Z. T., 845
Curkovic, T., 617
Cusick, D., 978

D

DaCosta, M. A., 723
Da Fonseca, C. R. V., 831
Dahlem, G. A., 218
Davis, D. R., 186
Decker, K. B., 300
De Coll, O., 70
DeHeer, C. J., 1101, 1213
de la Vega, M. H., 387
de León, J. H., 879
De Meyer, M., 261
Deyrup, L. D., 1199
Dmitriev, D. A., 187
Dole, S. A., 1034
Donahue, J. P., 723
Dorchin, N., 656
Dosdall, L. M., 64
Dossi, F. C. A., 1206
Duetting, P. S., 1146

E

Eastwood, R., 164
Edde, P. A., 175
Etxeberria, E., 1139
Evenden, M. L., 837

F

Farrell, B. D., 164
Favila, M. E., 1248
Feder, J., 662
Ferkovich, S. M., 586
Ferris, V. R., 241
Fordyce, J. A., 933
Forister, M. L., 933
Forrest, T. G., 978
Forschler, B. T., 1028
Fournier, F., 1121
Franklin, J., 536
Fraser, S., 1139
Freeman, T. P., 305
Fritz, A. H., 369
Fritz, G. N., 707

G

Galante, E., 1
Geils, B. W., 1133
Giblin-Davis, R. M., 352
Gibson, G., 799
Gillette, N. E., 433
Gold, R. E., 524
Gómez, B., 1
Gompert, Z., 933
González, R. H., 879
Gratton, C., 1182
Gribbins, K. M., 560
Gut, L. J., 898

H

Häfliger, P., 683
Hanks, L. M., 96, 718
Hardman, J. M., 536
Harris, M. O., 305
Hauser, M., 96
Havill, N. P., 195
Hayes, J. L., 945
Heinbockel, T., 988
Hendrichs, J., 387, 607
Henneberry, T. J., 917
Heraty, J. M., 567
Himuro, C., 974
Hirose, E., 91
Hirowatari, T., 1019
Ho, H.-Y., 859
Ho, J.-Z., 859
Hoddle, M. S., 553, 1154

CUMULATIVE AUTHOR INDEX
Annals of Entomology
Volume 99, 2006

A

Abrahamson, W. G., 656
Akimoto, S.-I., 457
Alarie, Y., 743
Albuquerque, G. S., 7
Aldrich, J. R., 91
Almeida, R. P. P., 577, 884
Aluja, M., 662
Amir-Maafi, M., 84
Amrine, J. W., Jr., 1057
Anderson, K. G., 305
Anderson, R. S., 1258, 1261
Archibald, S. B., 487
Ark, J. T., 560
Armbruster, P., 1234
Ashe, J. S., 19
Austin, J. W., 697

B

Barbour, J. D., 718
Barclay, H. J., 607
Beal, A. L. H., 545
Benoit, J. B., 560
Bentz, B. J., 189
Berlocher, S., 662
Birge, L. M., 449
Blossey, B., 683
Boivin, G., 1121
Bomar, C. R., 1257
Bostanian, N. J., 536
Boyce, S. L., 164
Boyd, E. A., 1154
Branch, C. L., 1146
Branham, M. A., 409
Brannon, S. L., 300
Braswell, W. E., 449
Brelsfoard, C. L., 707
Broce, A. B., 157
Brower, A. V. Z., 184
Brunner, J. F., 169, 617
Bugrov, A. G., 457
Burand, J. P., 969
Buschman, L. L., 279
Butler, C. D., 317
Byers, J. A., 865

C

Caccone, A., 195
Cáceres, C., 387
Caceres, C., 1191
Cagnotti, C. L., 70
Carmichael, A., 567
Carroll, M. K., 697
Castro, B. A., 999
Catarino, M., 1260
Cattelan, A. J., 91
Causton, C. E., 121
Cave, R. D., 776, 1146
Cavichioli, R. R., 648
Cayol, J.-P., 387
Cervantes, D. E., 718
Challet, G. L., 743
Chang, C. L., 1191
Chase, J. A., 463
Chatzimanolis, S., 19
Chen, T.-Y., 917
Cheng, J.-A., 1127
Chi, H., 84
Cho, I. K., 1191
Choi, M.-Y., 905
Christiansen, P., 342
Chu, C.-C., 917
Chuang, Y.-C., 859
Cisneros, R., 945
Cladera, J. L., 70
Coelho, J. R., 793
Cognato, A. I., 1034
Conn, J. E., 1234
Conte, H., 1206
Cook, W. M., 809
Copeland, R. S., 261, 530
Costa, C., 409
Cover, S. P., 487
Crampton, L., 33
Creel, O. R., 1146
Crespi, B. J., 407
Cruz, Z. T., 845
Curkovic, T., 617
Cusick, D., 978

D

DaCosta, M. A., 723
Da Fonseca, C. R. V., 831
Dahlem, G. A., 218
Davis, D. R., 186
Decker, K. B., 300
De Coll, O., 70
DeHeer, C. J., 1101, 1213
de la Vega, M. H., 387
de León, J. H., 879
De Meyer, M., 261
Deyrup, L. D., 1199
Dmitriev, D. A., 187
Dole, S. A., 1034
Donahue, J. P., 723
Dorchin, N., 656
Dosdall, L. M., 64
Dossi, F. C. A., 1206
Duetting, P. S., 1146

E

Eastwood, R., 164
Edde, P. A., 175
Etxeberria, E., 1139
Evenden, M. L., 837

F

Farrell, B. D., 164
Favila, M. E., 1248
Feder, J., 662
Ferkovich, S. M., 586
Ferris, V. R., 241
Fordyce, J. A., 933
Forister, M. L., 933
Forrest, T. G., 978
Forschler, B. T., 1028
Fournier, F., 1121
Franklin, J., 536
Fraser, S., 1139
Freeman, T. P., 305
Fritz, A. H., 369
Fritz, G. N., 707

G

Galante, E., 1
Geils, B. W., 1133
Giblin-Davis, R. M., 352
Gibson, G., 799
Gillette, N. E., 433
Gold, R. E., 524
Gómez, B., 1
Gompert, Z., 933
González, R. H., 879
Gratton, C., 1182
Gribbins, K. M., 560
Gut, L. J., 898

H

Häfliger, P., 683
Hanks, L. M., 96, 718
Hardman, J. M., 536
Harris, M. O., 305
Hauser, M., 96
Havill, N. P., 195
Hayes, J. L., 945
Heinbockel, T., 988
Hendrichs, J., 387, 607
Henneberry, T. J., 917
Heraty, J. M., 567
Himuro, C., 974
Hirose, E., 91
Hirowatari, T., 1019
Ho, H.-Y., 859
Ho, J.-Z., 859
Hoddle, M. S., 553, 1154

Hodgson, C. J., 121
Holliday, C. W., 793
Holt, R. D., 809
Hong, X.-Y., 959, 1057
Hoshizaki, S., 457
Hosokawa, T., 974
Hossler, E. W., 638
Hossler, F. E., 638
Howard, D. J., 449
Hsieh, C.-H., 768
Hsu, Y.-C., 859
Hu, C., 917
Hummel, N. A., 920

I

Ide, S., 409
Il'ichev, A. L., 898
Islam, A., 387

J

Jacobus, L. M., 241
Jang, E., 1191
Jiang, M.-X., 1127
Johnson, T. A., 938
Jones, V. P., 169
Joshi, R. C., 204
Juba, T. R., 1101
Judd, T. M., 398
Jurenka, R. A., 905

K

Kammlah, D. M., 966
Kasuya, E., 714, 1244
Kato, M., 851
Kawakita, A., 851
Kawano, K., 327
Keddie, B. A., 837
Kennedy, J. H., 821
Kim, J.-K., 1006
Kim, J.-L., 1006
Kirkendall, L. R., 211
Ko, C.-C., 768
Koji, S., 292
Křeček, J., 463
Krieger, M. J. B., 1213
Krupke, C. H., 169
Kubota, K., 1006
Kuriwada, T., 1244
Kuroko, H., 1019

L

Lacey, E. S., 718
Lachaud, J.-P., 567
Lajoie, D. R., 978
Landolt, P. J., 617
Landry, B., 121
Larson, P., 723
Lasnier, J., 536
Lawrence, J. F., 409
Lee, B.-W., 1019
Legaspi, J. C., 321
Lewis, J. L., 1028
Lewis, M. L., 204, 991
Lewter, J. A., 157
Liburd, O. E., 113
Linhart, Y. B., 1133
Liston, A., 33
Liu, C. Z., 1012
Lo, L.-W., 859
Logarzo, G. A., 879
Lohmeyer, K. H., 966
Lopez, M. S., 837
Lounibos, L. P., 1172
Luke, Q., 261
Lux, S., 261

M

Macchiavelli, R., 875
Machera, P., 261
Maharajh, B., 463
Mangold, J. R., 463
Mantoanelli, E., 7
Mathis, W. N., 999
Matsuura, K., 625
Matthews, R. W., 1199
McCafferty, W. P., 241
McKamey, S. H., 648
McKern, J., 697
McPherson, J. E., 755
Meikle, W. G., 253
Mendoza-López, M. R., 1248
Messenger, M. T., 697
Meyer, M. D., 241
Micó, E., 1
Miranda, X., 374
Miyatake, T., 364
Montgomery, M. E., 195
Moon, R., 799
Mooney, K. A., 1133
Moore, J. A., 305
Morales, M. A., 545
Moreau, C. S., 487
Moser, J. C., 891
Mu, J.-Y., 917
Muniappan, R., 845

N

Naczi, R.F.C., 218
Nakamura, K., 292
Nansen, C., 253
Nice, C. C., 933
Nigg, H. N., 1139

O

Ohba, S.-Y., 151
Oikawa, T., 364
Okumu, M., 261
O'Neil, R. J., 317
Ortíz-Domínguez, M., 1248
Overton, K., 524
Ovruski, S. M., 70
Owens, C. B., 157

P

Panizzi, A. R., 91
Pantoja, A., 875
Payne, S. A., 305
Peck, S. B., 121
Peng, C. Y.S., 920
Penny, N. D., 58
Pérez-Lachaud, G., 567
Perring, T. M., 598
Phillips, T. W., 175, 253
Pilon, J.-G., 1121
Polihronakis, M., 144
Poprawa, I., 592
Powell, J. A., 1257
Prado, S. S., 577
Pruett, J. H., 966
Purcell, A. H., 884

Q

Qureshi, J. A., 279

R

Rabaglia, R. J., 1034
Racette, G., 536
Radtke, M. G., 831
Ramaswamy, S. B., 279
Randolph, R. P., 241
Rao, A., 524
Rao, S., 33
Raulston, J. R., 1085
Reddy, G.V.P., 845
Rellinger, E. J., 560
Rinehart, J. E., 945
Rinehart, T. A., 113
Ripa, R., 352
Rivers, D. B., 1199
Robinson, A. S., 387
Rodríguez, R., 707
Roe, A. D., 433
Rohfritsch, O., 305
Roland, J., 1110
Roque-Albelo, L., 121
Ross, K. G., 1213
Rost, M. M., 592, 910
Rubinoff, D., 577
Rull, J., 662

S

Sabo, A., 241
Sakaluk, S. K., 989
Salazar, A., 875
Salinas-Moreno, Y., 945
Sampson, B. J., 113
Saran, R. K., 697
Sasakawa, K., 1006
Sasaki, M., 151
Scheffer, S. J., 204, 991
Scheffrahn, R. H., 352, 463
Schmidt, B. C., 1110
Schumann, R. A., 1139
Schwarzländer, M., 683
Scott, E. R., 656

Seal, J. N., 673
Segura, D. F., 70
Shapiro, A. M., 933
Shapiro, J. P., 321, 586
Shields, V. D. C., 988
Shima, H., 41
Shimamoto, K., 714
Shiyake, S., 195
Shoemaker, D. D., 1213
Shurtz, R. A., 755
Simmons, R., 185
Sinclair, B. J., 121
Six, D. L., 189
Smith, T. R., 776
Solodovnikov, A., 1064
Song, Z.-W., 1057
Souza, J. T. D., 91
Sperling, F. A. H., 433
Spiers, J. M., 113
Spurgeon, D. W., 1085
Stein, J. D., 433
Stelinski, L. L., 898
Stringer, S. J., 113
Stuart, R. J., 1139
Su, N.-Y., 352
Sun, L., 241
Suzuki, N., 974
Symmes, E. J., 598
Szalanski, A., 799
Szalanski, A. L., 157, 697

T

Tachi, T., 41
Takeyasu, J., 33
Takiya, D. M., 648
Tan, W., 969
Tatsuki, S., 457
Tatsuta, H., 151, 457
Tauber, C. A., 7
Tauber, M. J., 7
Taylor, D. B., 799
Taylor, J. M., 821
Thompson, V., 1259
Throne, J. E., 279
Tokuda, M., 629
Tooker, J. F., 96
Traoré, L., 1121
Triapitsyn, S. V., 879
Tripodi, A. D., 697
Tschinkel, W. R., 673
Tsuji, K., 851

U

Uechi, N., 1165
Ulmer, B. J., 64

V

Vaccaro, N., 70
Vanin, S. A., 409
Vargas, R. I., 1191
Vargo, E. L., 1101
Vera, M. T., 70, 387
Vinson, S. B., 524
Virla, E. G., 879
Voss, S. C., 755
Vreysen, M. J. B., 607

W

Wagner, D. L., 638
Wang, C.-H., 768
Wang, G., 1012
Warchalowska-Sliwa, E., 457
Way, M. O., 999, 1127
Weaver, S. C., 1172
Weller, S. J., 723
Wharton, R. A., 261, 530
Williamson, G. B., 831
Williamson, R. C., 938
Wornoayporn, V., 387

X

Xie, L., 959
Xue, X.-F., 959, 1057

Y

Yan, L., 1012
Yanagi, S.-I., 364
Yang, J. J., 1139
Yanoviak, S. P., 1172
Yasumoto, A. A., 714
Ye, W., 352
Yeargan, K. V., 300
Yoder, J. A., 560
Yoder, R., 1127
Yu, G., 195
Yukawa, J., 629, 1165

Z

Zacaro, A. A., 1206
Zalom, F. G., 920
Zatwarnicki, T., 999
Zenz, N., 261
Zhang, W.-J., 1127
Zúñiga, G., 945

CUMULATIVE KEY WORD INDEX
Annals of Entomology
Volume 99, 2006

A

abundance,
  bamboo, 1172
  deforestation, 1172
  diversity, 1172
  tree hole, 1172
acoustic behavior,
  communication, 978
  phonoresponse, 978
Acronicta,
  larval evolution, 638
  Nyssa, 638
  wood-boring, 638
adult morphology,
  androconia, 723
  phylogeny, 723
  ultrasound, 723
Aedes albopictus,
  invasion biology, 1234
  larval growth, 1234
  life history, 1234
Aedes vexans,
  genetic variation, 157
  mitochondrial DNA, 157
  population structure, 157
aggregation,
  diel activity patterns, 169
  Euschistus conspersus, 169
  feeding, 169
  mating, 169
agricultural ecosystems,
  crane fly pest species, 33
  DNA diagnosis, 33
  Oregon, 33
  urban landscapes, 33
Agromyzidae, 204, 991
  cryptic species, 991
  DNA barcodes, 204
  DNA taxonomy, 204
  host-race, 991
  invasive species, 991
  leafminer, 991
  Liriomyza, 204
alate, 524, 625
  developmental constraint, 625
  eclosion, 625
  imported fire ant, 524
  infection, 524
  microbial control, 524
  pathogen, 524
  sexual dimorphism, 625
  swarming, 625
Albitarsis Complex,
  Anopheles marajoara, 707
  distribution, 707
  ITS2, 707
  taxonomy, 707
Aleyrodidae,
  Asia, 768
  Bemisia tabaci, 768
  mitochondrial DNA cytochrome oxidase I, 768
  molecular marker, 768
alkenal,
  dorsal abdominal gland, 859
  exuviae, 859
  limonene, 859
  pinene, 859
allometry, 327, 342
  body size, 327
  intraspecific variability, 327
  Megasoma, 342
  oversized apparatus, 327
  phylogenetic inertia, 327
  sexual selection, 342
  somatic proportions, 342
Allonemobius,
  calling song, 449
  Nemobiinae, 449
  Orthoptera, 449
  phylogeny, 449
alternative mating strategy,
  body size, 974
  courtship behavior, 974
  mating success, 974
  Megacopta punctatissima, 974
ambrosia beetles,
  exotic species, 1034
  invasive pests, 1034
  Scolytidae, 1034
  Xyleborini, 1034
anaerobic protists identification,
  Oxymonadida, 1028
  Parabasalia, 1028
  symbiotic protists, 1028
  termite identification, 1028
Anastrepha suspensa,
  5.8S, 369
  ITS2, 369
  rDNA, 369
  Tephritidae, 369
Andex insignis,
  Dytiscidae, 743
  Hyphydrini, 743
  larvae, 743
  phylogeny, 743
androconia,
  adult morphology, 723
  phylogeny, 723
  ultrasound, 723
Anopheles marajoara,
  Albitarsis Complex, 707
  distribution, 707
  ITS2, 707
  taxonomy, 707
ant, 398, 545
  food storage, 398
  foraging, 398
  Membracidae, 545
  mutualism, 545
  Pheidole ceres, 398
  seeds, 398
  Solidago altissima, 545
ant evolution,
  Eocene, 487
  Formicidae, 487
  Myrmeciinae, 487
  Okanagan Highlands, 487
Anthonomus grandis,
  boll weevil, 1085
  photoperiod, 1085
  reproductive diapause, 1085
Antispila distyliella,
  Heliozelidae, 1019
  immature stage, 1019
  Japan, 1019
  Lepidoptera, 1019
ant–host association,
  Ectatomma, 567
  Eucharitidae, 567
  Kapala, 567
  Ponerinae, 567
Anystidae,
  biological control, 536
  Erythraeidae, 536
  Phytoseiidae, 536
  Stigmaeidae, 536
Asia,
  Aleyrodidae, 768
  Bemisia tabaci, 768
  mitochondrial DNA cytochrome oxidase I, 768
  molecular marker, 768
Asphondylia,
  Cecidomyiidae, 1165
  host range, 1165
  life history strategy, 1165
  short-term alternative hosts, 1165
Asphondyliini,
  Bruggmanniella, 629
  Cecidomyiidae, 629
  gall midge, 629

Lauraceae, 629
asymmetry,
development, 7
larvae, 7
Leucochrysinae, 7
phenotype, 7
Atta texana,
central cavity, 891
feeder holes, 891
fungus gardens, 891
nest excavation, 891
attachment,
Graphocephala, 884
Homalodisca, 884
Pierce's disease, 884
Attini,
Florida, 673
fungus gardening, 673
longleaf pine, 673
sandhills, 673
autodetection,
female response, 898
Grapholita molesta, 898
Oriental fruit moth, 898
sex pheromone, 898

B

bacteria,
Chrysoperla carnea, 917
larvae, 917
midgut, 917
yeast, 917
bait design,
bait station, 1139
Caribbean fruit fly, 1139
bait station,
bait design, 1139
Caribbean fruit fly, 1139
Balaustium,
moisture requirements, 560
velvet mite, 560
water vapor uptake, 560
bamboo, 1057, 1172
abundance, 1172
deforestation, 1172
diversity, 1172
eriophyids, 1057
key, 1057
Qinling Mountains, 1057
taxonomy, 1057
tree hole, 1172
bark beetle, 189, 945
Dendroctonus, 945
genetic structure, 945
isozyme, 945
Mexico, 945
mycangial fungi, 189
mycophagy, 189
symbiosis, 189
bee,
body size, 714
Colletes perforator, 714
male, 714
mating, 714
beetles,
Cybocephalus, 776
revision, 776
scale predator, 776
taxonomy, 776
behavior, 253, 598
Bemisia tabaci biotype A, 598
egg distribution, 253
ovipositional strategy, 253
stored-products, 253
Trialeurodes vaporariorum, 598
whitefly, 598
Bemisia tabaci,
Aleyrodidae, 768
Asia, 768
mitochondrial DNA cytochrome oxidase I, 768
molecular marker, 768
Bemisia tabaci biotype A,
behavior, 598
Trialeurodes vaporariorum, 598
whitefly, 598
bigeyed bug,
critical photoperiod, 300
overwintering, 300
sensitive stage, 300
biocontrol,
Cecidochares, 845
Chromolaena, 845
gall, 845
nutrient sink, 845
biogeography, 1, 463, 933
diversity survey, 463
identification keys, 463
Mesoamerica, 1
new species, 1, 463
Remington suture zone, 933
Rutelidae, 1
scanning electron microscopy, 933
specialization, 933
West Indies, 463
Yaaxkumukia, 1
biological control, 96, 113, 536, 879
Anystidae, 536
Charles Robertson, 96
Erythraeidae, 536
Eulophidae, 113
floral resources, 96
fly, 96
Gonatocerus tuberculifemur, 879
high-fidelity polymerase chain reaction, 113
Homalodisca coagulata, 879
parasitism, 113
Phytoseiidae, 536
Pierce's disease, 879
Platygastridae, 113
pollinator, 96
Stigmaeidae, 536
biological weed control,
common reed, 683
life history, 683
stem borer, 683
wetlands, 683
biomass,
Neotropical Scarabaeidae, 831
preservatives, 831
volume, 831
volume–mass relationship, 831
body size, 211, 327, 714, 837, 974
allometry, 327
alternative mating strategy, 974
bee, 714
Colletes perforator, 714
Costa Rica, 211
courtship behavior, 974
host specificity, 211
intraspecific variability, 327
large aspen tortrix, 837
male, 714
mating, 714
mating success, 974
Megacopta punctatissima, 974
microsporidia, 837
oversized apparatus, 327
phylogenetic inertia, 327
reproductive performance, 837
Scolytinae, 211
Xyleborini, 211
boll weevil,
Anthonomus grandis, 1085
photoperiod, 1085
reproductive diapause, 1085
boreal forest,
community composition, 1110
landscape structure, 1110
Malacosoma disstria, 1110
parasitoid, 1110
Brachypsectra,
fossil new species, 409
identification key, 409
Miocene amber, 409
relationships, 409
bromeliads,
development, 1146
fecundity, 1146
Metamasius callizona, 1146
weevil, 1146
Bruggmanniella,
Asphondyliini, 629
Cecidomyiidae, 629
gall midge, 629
Lauraceae, 629
Bt corn,
insect dispersal, 279
resistance management, 279
southwestern corn borer, 279

transgenic crops, 279
Buchnera,
Heteroptera, 577
maximum likelihood, 577
stinkbug, 577
symbiosis, 577
buck moth,
Hemileuca maia species complex, 1182
host shift, 1182
novel host, 1182
preference–performance, 1182
bulking agent,
liquid diet rearing, 1191
oriental fruit fly, 1191
sponge cloth, 1191

C

Caenidae,
Ephemeroptera, 821
life history, 821
secondary production, 821
calcium channel,
PBAN, 905
pheromone biosynthesis, 905
receptor signal transduction, 905
Cales noacki,
demography, 553
fortuitous biological control, 553
jackknife, 553
Persea americana, 553
calling song,
Allonemobius, 449
Nemobiinae, 449
Orthoptera, 449
phylogeny, 449
Callosobruchus chinensis,
egg number, 364
egg size, 364
genetic correlation, 364
trade-off, 364
canola,
Ceutorhynchus obstrictus, 64
emergence, 64
monitoring, 64
sex ratio, 64
Canthon cyanellus,
ecological speciation, 1248
prezygotic isolation, 1248
sexual recognition, 1248
Caribbean fruit fly,
bait design, 1139
bait station, 1139
Cassida rubiginosa,
demographic traits, 292
mark–recapture, 292
population stability, 292
reproductive rate, 292
Cecidochares,
biocontrol, 845
Chromolaena, 845
gall, 845
nutrient sink, 845
Cecidomyiidae, 629, 1165
Asphondylia, 1165
Asphondyliini, 629
Bruggmanniella, 629
gall midge, 629
host range, 1165
Lauraceae, 629
life history strategy, 1165
short-term alternative hosts, 1165
cell wall,
host–parasite interface, 305
plasmodesmata, 305
sink tissues, 305
transmission electron micrograph, 305
cellularization,
cleavage, 592
embryogenesis, 592
energids, 592
midgut, 592
central cavity,
Atta texana, 891
feeder holes, 891
fungus gardens, 891
nest excavation, 891
Ceratitis anonae,
Ceratitis fasciventris, 530
Ceratitis rosa, 530
Solanaceae, 530
Ceratitis fasciventris,
Ceratitis anonae, 530
Ceratitis rosa, 530
Solanaceae, 530
Ceratitis rosa,
Ceratitis anonae, 530
Ceratitis fasciventris, 530
Solanaceae, 530
Ceutorhynchus obstrictus,
canola, 64
emergence, 64
monitoring, 64
sex ratio, 64
Charles Robertson,
biological control, 96
floral resources, 96
fly, 96
pollinator, 96
Choristoneura rosaceana,
courtship behavior, 617
Pandemis pyrusana, 617
Chromolaena,
biocontrol, 845
Cecidochares, 845
gall, 845
nutrient sink, 845
chromosome rearrangement,
introgression, 457
mtDNA, 457
phylogeography, 457
Robertsonian fusion, 457
Chrysoperla carnea,
bacteria, 917
larvae, 917
midgut, 917
yeast, 917
Chrysopidae,
key, 58
Meleoma, 58
Neuroptera, 58
taxonomy, 58
cicada killer,
cicada-killer, 793
key, 793
New World, 793
Sphecius, 793
cicadas,
colonization, 809
heterogeneous habitats, 809
host plant density, 809
rough-leaved dogwood, 809
citrus pest,
genetic markers, 164
Hispaniola, 164
invasive species, 164
West Indies, 164
cladistic analyses,
endophallus, 1006
phylogeny, 1006
Pristosia impunctata sp. nov., 1006
Pristosia latistoma sp. nov., 1006
cleavage,
cellularization, 592
embryogenesis, 592
energids, 592
midgut, 592
COII,
12S, 352
16S, 352
molecular phylogenetics, 352
phylogenetics, 352
Colletes perforator,
bee, 714
body size, 714
male, 714
mating, 714
colonization,
cicadas, 809
heterogeneous habitats, 809
host plant density, 809
rough-leaved dogwood, 809
colony genetic structure,
eastern subterranean termite, 1101
Formosan subterranean termite, 1101

microsatellite, 1101
population genetics, 1101
color analysis,
cotton, 865
image analysis, 865
RGB, 865
vision, 865
color dimorphism,
elytra, 1127
Lissorhoptrus oryzophilus, 1127
mating, 1127
ovarian development, 1127
common reed,
biological weed control, 683
life history, 683
stem borer, 683
wetlands, 683
communication,
acoustic behavior, 978
phonoresponse, 978
community composition,
boreal forest, 1110
landscape structure, 1110
Malacosoma disstria, 1110
parasitoid, 1110
community ecology,
detritus, 1133
indirect effect, 1133
plant morphology, 1133
tritrophic, 1133
Cosmopolites sordidus,
head capsules width, 875
instars, 875
plantains, 875
precipitation, 875
Costa Rica,
body size, 211
host specificity, 211
Scolytinae, 211
Xyleborini, 211
cotton,
color analysis, 865
image analysis, 865
RGB, 865
vision, 865
courtship behavior, 617, 974
alternative mating strategy, 974
body size, 974
Choristoneura rosaceana, 617
mating success, 974
Megacopta punctatissima, 974
Pandemis pyrusana, 617
crane fly pest species,
agricultural ecosystems, 33
DNA diagnosis, 33
Oregon, 33
urban landscapes, 33
cricket,
life history strategy, 1244
mate choice, 1244
sexual selection, 1244
critical photoperiod,
bigeyed bug, 300
overwintering, 300
sensitive stage, 300
cryptic species,
Agromyzidae, 991
host-race, 991
invasive species, 991
leafminer, 991
cryptic species complex,
reproductive compatibility, 387
sexual isolation, 387
sterile insect technique, 387
temporal isolation, 387
cultivated,
Fopius, 261
indigenous, 261
Opiinae, 261
Psyttalia, 261
Cybocephalus,
beetles, 776
revision, 776
scale predator, 776
taxonomy, 776
cytochrome oxidase *c*,
Dioryctria, 433
DNA barcoding, 433
morphology, 433
mtDNA, 433
cytochrome oxidase subunit I,
mitochondrial DNA, 959
phylogenetic analysis, 959
Tetranychus urticae, 959

D

deforestation,
abundance, 1172
bamboo, 1172
diversity, 1172
tree hole, 1172
demographic traits,
Cassida rubiginosa, 292
mark–recapture, 292
population stability, 292
reproductive rate, 292
demography,
Cales noacki, 553
fortuitous biological control, 553
jackknife, 553
Persea americana, 553
Dendroctonus,
bark beetle, 945
genetic structure, 945
isozyme, 945
Mexico, 945
detritus,
community ecology, 1133
indirect effect, 1133
plant morphology, 1133
tritrophic, 1133
development, 7, 1146
asymmetry, 7
bromeliads, 1146
fecundity, 1146
larvae, 7
Leucochrysinae, 7
Metamasius callizona, 1146
phenotype, 7
weevil, 1146
developmental constraint,
alate, 625
eclosion, 625
sexual dimorphism, 625
swarming, 625
developmental delay,
Melittobia digitata, 1199
paralysis, 1199
venom, 1199
Diatraea,
embryonic stages, 1206
organogenesis, 1206
RNA levels, 1206
diel activity patterns,
aggregation, 169
Euschistus conspersus, 169
feeding, 169
mating, 169
Dioryctria,
cytochrome oxidase *c*, 433
DNA barcoding, 433
morphology, 433
mtDNA, 433
Diptera,
distribution, 662
host plant, 662
Rhagoletis, 662
Tephritidae, 662
dispersal,
mate choice, 607
remating, 607
sperm, 607
sterile insect technique, 607
distribution, 662, 707
Albitarsis Complex, 707
Anopheles marajoara, 707
Diptera, 662
host plant, 662
ITS2, 707
Rhagoletis, 662
taxonomy, 707
Tephritidae, 662
diversity,
abundance, 1172
bamboo, 1172
deforestation, 1172
tree hole, 1172
diversity survey,
biogeography, 463
identification keys, 463
new species, 463
West Indies, 463

DNA barcodes,
Agromyzidae, 204
DNA taxonomy, 204
Liriomyza, 204
DNA barcoding,
cytochrome oxidase *c*, 433
Dioryctria, 433
morphology, 433
mtDNA, 433
DNA diagnosis,
agricultural ecosystems, 33
crane fly pest species, 33
Oregon, 33
urban landscapes, 33
DNA taxonomy,
Agromyzidae, 204
DNA barcodes, 204
Liriomyza, 204
dorsal abdominal gland,
alkenal, 859
exuviae, 859
limonene, 859
pinene, 859
Dyar's rule,
instar determination, 938
linden borer, 938
Dytiscidae,
Andex insignis, 743
Hyphydrini, 743
larvae, 743
phylogeny, 743

E

18s rDNA,
Ephemeroptera, 241
phylogeny, 241
eastern subterranean termite,
colony genetic structure, 1101
Formosan subterranean termite, 1101
microsatellite, 1101
population genetics, 1101
eclosion,
alate, 625
developmental constraint, 625
sexual dimorphism, 625
swarming, 625
ecological speciation,
Canthon cyanellus, 1248
prezygotic isolation, 1248
sexual recognition, 1248
Ectatomma,
ant–host association, 567
Eucharitidae, 567
Kapala, 567
Ponerinae, 567
egg,
ELISA, 586
oviposition, 586
Parafilm, 586
predator, 586
egg distribution,
behavior, 253
ovipositional strategy, 253
stored-products, 253
egg number,
Callosobruchus chinensis, 364
egg size, 364
genetic correlation, 364
trade-off, 364
egg size,
Callosobruchus chinensis, 364
egg number, 364
genetic correlation, 364
trade-off, 364
ELISA, 321, 586
eggs, 586
food quality, 321
nutrition, 321
oviposition, 586
Parafilm, 586
predation, 321
predator, 586
reproduction, 321
elytra,
color dimorphism, 1127
Lissorhoptrus oryzophilus, 1127
mating, 1127
ovarian development, 1127
embryogenesis,
cellularization, 592
cleavage, 592
energids, 592
midgut, 592
embryonic stages,
Diatraea, 1206
organogenesis, 1206
RNA levels, 1206
emergence,
canola, 64
Ceutorhynchus obstrictus, 64
monitoring, 64
sex ratio, 64
endophallus,
cladistic analyses, 1006
phylogeny, 1006
Pristosia impunctata sp. nov., 1006
Pristosia latistoma sp. nov., 1006
energids,
cellularization, 592
cleavage, 592
embryogenesis, 592
midgut, 592
Ennya,
jamming, 374
Membracidae, 374
substrate-borne vibrations, 374
enteric bacteria,
Enterococcus faecalis, 91
Klebsiella pneumoniae, 91
Nezara viridula, 91
Enterococcus faecalis,
enteric bacteria, 91
Klebsiella pneumoniae, 91
Nezara viridula, 91
Eocene,
ant evolution, 487
Formicidae, 487
Myrmeciinae, 487
Okanagan Highlands, 487
Ephemeroptera, 241, 821
18s rDNA, 241
Caenidae, 821
life history, 821
phylogeny, 241
secondary production, 821
eriophyids,
bamboo, 1057
key, 1057
Qinling Mountains, 1057
taxonomy, 1057
Erythraeidae,
Anystidae, 536
biological control, 536
Phytoseiidae, 536
Stigmaeidae, 536
Eucharitidae,
ant–host association, 567
Ectatomma, 567
Kapala, 567
Ponerinae, 567
Eulophidae,
biological control, 113
high-fidelity polymerase chain reaction, 113
parasitism, 113
Platygastridae, 113
Euschistus conspersus,
aggregation, 169
diel activity patterns, 169
feeding, 169
mating, 169
exotic species,
ambrosia beetles, 1034
invasive pests, 1034
Scolytidae, 1034
Xyleborini, 1034
exuviae,
alkenal, 859
dorsal abdominal gland, 859
limonene, 859
pinene, 859
eye color,
Hematobia irritans, 966
mutant, 966

F

fecundity,
bromeliads, 1146
development, 1146
Metamasius callizona, 1146
weevil, 1146

feeder holes,
Atta texana, 891
central cavity, 891
fungus gardens, 891
nest excavation, 891
feeding,
aggregation, 169
diel activity patterns, 169
Euschistus conspersus, 169
mating, 169
female response,
autodetection, 898
Grapholita molesta, 898
Oriental fruit moth, 898
sex pheromone, 898
filth fly parasitoids,
molecular diagnostics, 799
Pteromalidae, 799
Spalangia, 799
fire ants,
genetic structure, 1213
invasion history, 1213
Solenopsis invicta, 1213
Fletcherimyia,
pitcher plants, 218
Sarcophagidae, 218
Sarraceniaceae, 218
systematics, 218
floral resources,
biological control, 96
Charles Robertson, 96
fly, 96
pollinator, 96
Florida,
Attini, 673
fungus gardening, 673
longleaf pine, 673
sandhills, 673
fly,
biological control, 96
Charles Robertson, 96
floral resources, 96
pollinator, 96
food quality,
ELISA, 321
nutrition, 321
predation, 321
reproduction, 321
food storage,
ants, 398
foraging, 398
Pheidole ceres, 398
seeds, 398
Fopius,
cultivated, 261
indigenous, 261
Opiinae, 261
Psyttalia, 261
foraging,
ants, 398
food storage, 398
Pheidole ceres, 398
seeds, 398
Formicidae,
ant evolution, 487
Eocene, 487
Myrmeciinae, 487
Okanagan Highlands, 487
Formosan subterranean termite,
colony genetic structure, 1101
eastern subterranean termite, 1101
microsatellite, 1101
population genetics, 1101
fortuitous biological control,
Cales noacki, 553
demography, 553
jackknife, 553
Persea americana, 553
fossil new species,
Brachypsectra, 409
identification key, 409
Miocene amber, 409
relationships, 409
fungus gardening,
Attini, 673
Florida, 673
longleaf pine, 673
sandhills, 673
fungus gardens,
Atta texana, 891
central cavity, 891
feeder holes, 891
nest excavation, 891

G

gall,
biocontrol, 845
Cecidochares, 845
Chromolaena, 845
nutrient sink, 845
gall midge, 629, 656
Asphondyliini, 629
Bruggmanniella, 629
Cecidomyiidae, 629
inquiline, 656
Lauraceae, 629
Solidago, 656
Gastrodia,
mycoheterotrophy, 851
orchid, 851
pollination, 851
Scatophagidae, 851
genetic correlation,
Callosobruchus chinensis, 364
egg number, 364
egg size, 364
trade-off, 364
genetic markers,
citrus pest, 164
Hispaniola, 164
invasive species, 164
West Indies, 164
genetic structure, 945, 1213
bark beetle, 945
Dendroctonus, 945
fire ants, 1213
invasion history, 1213
isozyme, 945
Mexico, 945
Solenopsis invicta, 1213
genetic variation, 157, 697
Aedes vexans, 157
mitochondrial DNA, 157, 697
population structure, 157
Reticulitermes, 697
termite, 697
genitalic evolution,
intraspecific variation, 144
morphometrics, 144
Phyllophaga hirticula, 144
postcopulatory sexual selection, 144
geographical distribution,
Mediterranean fruit fly, 70
relative abundance index, 70
Tephritidae, 70
glassy-winged sharpshooter, 648, 920
Homalodisca coagulata, 648
leafhopper, 920
oogenesis, 920
Pierce's disease, 648
Propetes, 648
sperm, 920
telotrophic, 920
Xylella fastidiosa, 648
gonad-specific virus,
Helicoverpa zea, 969
Hz-2V, 969
mating behavior, 969
pheromonostatic peptide, 969
Gonatocerus latipennis,
leafhopper biology, 1154
phenology, 1154
Polynema sp., 1154
Vitis girdiana, 1154
Gonatocerus tuberculifemur,
biological control, 879
Homalodisca coagulata, 879
Pierce's disease, 879
Graphocephala,
attachment, 884
Homalodisca, 884
Pierce's disease, 884
Grapholita molesta,
autodetection, 898
female response, 898
Oriental fruit moth, 898
sex pheromone, 898
growth trajectory,
Lethocerus deyrolli, 151
morphology, 151
phenotypic plasticity, 151
predation, 151

Gynaephora menyuanensis,
head capsule width, 1012
instars, 1012
life cycle, 1012
stadium duration, 1012

H

head capsule width, 875, 1012
Cosmopolites sordidus, 875
Gynaephora menyuanensis, 1012
instars, 875, 1012
life cycle, 1012
plantains, 875
precipitation, 875
stadium duration, 1012
Helicoverpa zea,
gonad-specific virus, 969
Hz-2V, 969
mating behavior, 969
pheromonostatic peptide, 969
Heliozelidae,
Antispila distyliella, 1019
immature stage, 1019
Japan, 1019
Lepidoptera, 1019
Hematobia irritans,
eye color, 966
mutant, 966
Hemileuca maia species complex,
buck moth, 1182
host shift, 1182
novel host, 1182
preference–performance, 1182
heterogeneous habitats,
cicadas, 809
colonization, 809
host plant density, 809
rough-leaved dogwood, 809
Heteroptera,
Buchnera, 577
maximum likelihood, 577
stinkbug, 577
symbiosis, 577
high-fidelity polymerase chain reaction,
biological control, 113
Eulophidae, 113
parasitism, 113
Platygastridae, 113
Hispaniola,
citrus pest, 164
genetic markers, 164
invasive species, 164
West Indies, 164
Homalodisca,
attachment, 884
Graphocephala, 884
Pierce's disease, 884
Homalodisca coagulata, 648, 879
biological control, 879
glassy-winged sharpshooter, 648
Gonatocerus tuberculifemur, 879
Pierce's disease, 648, 879
Propetes, 648
Xylella fastidiosa, 648
host plant,
Diptera, 662
distribution, 662
Rhagoletis, 662
Tephritidae, 662
host plant density,
cicadas, 809
colonization, 809
heterogeneous habitats, 809
rough-leaved dogwood, 809
host range,
Asphondylia, 1165
Cecidomyiidae, 1165
life history strategy, 1165
short-term alternative hosts, 1165
host shift,
buck moth, 1182
Hemileuca maia species complex, 1182
novel host, 1182
preference–performance, 1182
host specificity,
body size, 211
Costa Rica, 211
Scolytinae, 211
Xyleborini, 211
host-race,
Agromyzidae, 991
cryptic species, 991
invasive species, 991
leafminer, 991
host–parasite interface,
cell wall, 305
plasmodesmata, 305
sink tissues, 305
transmission electron micrograph, 305
Hydrellia wirthi Korytkowski,
invasive species, 999
shore flies, 999
Hyphydrini,
Andex insignis, 743
Dytiscidae, 743
larvae, 743
phylogeny, 743
Hz-2V,
gonad-specific virus, 969
Helicoverpa zea, 969
mating behavior, 969
pheromonostatic peptide, 969

I

identification key, 409, 463
biogeography, 463
Brachypsectra, 409
diversity survey, 463
fossil new species, 409
Miocene amber, 409
new species, 463
relationships, 409
West Indies, 463
image analysis,
color analysis, 865
cotton, 865
RGB, 865
vision, 865
immature stage,
Antispila distyliella, 1019
Heliozelidae, 1019
Japan, 1019
Lepidoptera, 1019
impact,
introduced species, 121
invasive, 121
prioritization, 121
imported fire ant,
alate, 524
infection, 524
microbial control, 524
pathogen, 524
indigenous,
cultivated, 261
Fopius, 261
Opiinae, 261
Psyttalia, 261
indirect effect,
community ecology, 1133
detritus, 1133
plant morphology, 1133
tritrophic, 1133
infection,
alate, 524
imported fire ant, 524
microbial control, 524
pathogen, 524
inquiline,
gall midges, 656
Solidago, 656
insect dispersal,
Bt corn, 279
resistance management, 279
southwestern corn borer, 279
transgenic crops, 279
instar determination,
Dyar's rule, 938
linden borer, 938
instars, 755, 875, 1012
Cosmopolites sordidus, 875
Gynaephora menyuanensis, 1012
head capsule width, 1012
head capsules width, 875
life cycle, 1012

midwestern species, 755
nymphal key, 755
plantains, 875
precipitation, 875
Sinea, 755
stadium duration, 1012
intraspecific variability,
allometry, 327
body size, 327
oversized apparatus, 327
phylogenetic inertia, 327
intraspecific variation,
genitalic evolution, 144
morphometrics, 144
Phyllophaga hirticula, 144
postcopulatory sexual selection, 144
introduced species,
impact, 121
invasive, 121
prioritization, 121
introgression,
chromosome rearrangement, 457
mtDNA, 457
phylogeography, 457
Robertsonian fusion, 457
invasion biology,
Aedes albopictus, 1234
larval growth, 1234
life history, 1234
invasion history,
fire ants, 1213
genetic structure, 1213
Solenopsis invicta, 1213
invasive,
impact, 121
introduced species, 121
prioritization, 121
invasive pest, 195, 1034
ambrosia beetles, 1034
exotic species, 1034
molecular systematics, 195
Scolytidae, 1034
Tsuga, 195
Xyleborini, 1034
invasive species, 164, 991, 999
Agromyzidae, 991
citrus pest, 164
cryptic species, 991
genetic markers, 164
Hispaniola, 164
host-race, 991
Hydrellia wirthi Korytkowski, 999
leafminer, 991
shore flies, 999
West Indies, 164
isozyme,
bark beetle, 945
Dendroctonus, 945
genetic structure, 945
Mexico, 945
ITS2, 369, 707
5.8S, 369
Albitarsis Complex, 707
Anastrepha suspensa, 369
Anopheles marajoara, 707
distribution, 707
rDNA, 369
taxonomy, 707
Tephritidae, 369

J

jackknife,
Cales noacki, 553
demography, 553
fortuitous biological control, 553
Persea americana, 553
jamming,
Ennya, 374
Membracidae, 374
substrate-borne vibrations, 374
Japan,
Antispila distyliella, 1019
Heliozelidae, 1019
immature stage, 1019
Lepidoptera, 1019

K

Kapala,
ant–host association, 567
Ectatomma, 567
Eucharitidae, 567
Ponerinae, 567
key, 58, 793, 1057
bamboo, 1057
Chrysopidae, 58
eriophyids, 1057
Meleoma, 58
Neuroptera, 58
New World, 793
Qinling Mountains, 1057
Sphecius, 793
taxonomy, 58, 1057
Klebsiella pneumoniae,
enteric bacteria, 91
Enterococcus faecalis, 91
Nezara viridula, 91

L

landscape structure,
boreal forest, 1110
community composition, 1110
Malacosoma disstria, 1110
parasitoid, 1110
large aspen tortrix,
body size, 837
microsporidia, 837
reproductive performance, 837
larvae, 7, 743, 917
Andex insignis, 743
asymmetry, 7
bacteria, 917
Chrysoperla carnea, 917
development, 7
Dytiscidae, 743
Hyphydrini, 743
Leucochrysinae, 7
midgut, 917
phenotype, 7
phylogeny, 743
yeast, 917
larval evolution,
Acronicta, 638
Nyssa, 638
wood-boring, 638
larval growth,
Aedes albopictus, 1234
invasion biology, 1234
life history, 1234
Lauraceae,
Asphondyliini, 629
Bruggmanniella, 629
Cecidomyiidae, 629
gall midge, 629
leafhopper,
glassy-winged sharpshooter, 920
oogenesis, 920
sperm, 920
telotrophic, 920
leafhopper biology,
Gonatocerus latipennis, 1154
phenology, 1154
Polynema sp., 1154
Vitis girdiana, 1154
leafminer,
Agromyzidae, 991
cryptic species, 991
host-race, 991
invasive species, 991
Lepidoptera,
Antispila distyliella, 1019
Heliozelidae, 1019
immature stage, 1019
Japan, 1019
Lethocerus deyrolli,
growth trajectory, 151
morphology, 151
phenotypic plasticity, 151
predation, 151
Leucochrysinae,
asymmetry, 7
development, 7
larvae, 7
phenotype, 7
life cycle,
Gynaephora menyuanensis, 1012
head capsule width, 1012
instars, 1012
stadium duration, 1012

life history, 683, 821, 1234
Aedes albopictus, 1234
biological weed control, 683
Caenidae, 821
common reed, 683
Ephemeroptera, 821
invasion biology, 1234
larval growth, 1234
secondary production, 821
stem borer, 683
wetlands, 683
life history strategy, 1165, 1244
Asphondylia, 1165
Cecidomyiidae, 1165
cricket, 1244
host range, 1165
mate choice, 1244
sexual selection, 1244
short-term alternative hosts, 1165
life table,
parasitoid, 84
reproduction, 84
stored product pest, 84
limonene,
alkenal, 859
dorsal abdominal gland, 859
exuviae, 859
pinene, 859
linden borer,
Dyar's rule, 938
instar determination, 938
liquid diet rearing,
bulking agent, 1191
oriental fruit fly, 1191
sponge cloth, 1191
Liriomyza,
Agromyzidae, 204
DNA barcodes, 204
DNA taxonomy, 204
Lissorhoptrus oryzophilus,
color dimorphism, 1127
elytra, 1127
mating, 1127
ovarian development, 1127
longleaf pine,
Attini, 673
Florida, 673
fungus gardening, 673
sandhills, 673

M

Malacosoma disstria,
boreal forest, 1110
community composition, 1110
landscape structure, 1110
parasitoid, 1110
male,
bee, 714
body size, 714
Colletes perforator, 714
mating, 714
mark–recapture,
Cassida rubiginosa, 292
demographic traits, 292
population stability, 292
reproductive rate, 292
mate choice, 607, 1244
cricket, 1244
dispersal, 607
life history strategy, 1244
remating, 607
sexual selection, 1244
sperm, 607
sterile insect technique, 607
mate location,
Prionus californicus, 718
semiochemical, 718
wood borer, 718
mating, 169, 714, 1127
aggregation, 169
bee, 714
body size, 714
Colletes perforator, 714
color dimorphism, 1127
diel activity patterns, 169
elytra, 1127
Euschistus conspersus, 169
feeding, 169
Lissorhoptrus oryzophilus, 1127
male, 714
ovarian development, 1127
mating behavior,
gonad-specific virus, 969
Helicoverpa zea, 969
Hz-2V, 969
pheromonostatic peptide, 969
mating success,
alternative mating strategy, 974
body size, 974
courtship behavior, 974
Megacopta punctatissima, 974
maximum likelihood,
Buchnera, 577
Heteroptera, 577
stinkbug, 577
symbiosis, 577
Mediterranean fruit fly,
geographical distribution, 70
relative abundance index, 70
Tephritidae, 70
Megacopta punctatissima,
alternative mating strategy, 974
body size, 974
courtship behavior, 974
mating success, 974
Megasoma,
allometry, 342
sexual selection, 342
somatic proportions, 342
Meleoma,
Chrysopidae, 58
key, 58
Neuroptera, 58
taxonomy, 58
Melittobia digitata,
developmental delay, 1199
paralysis, 1199
venom, 1199
Membracidae, 374, 545
ants, 545
Ennya, 374
jamming, 374
mutualism, 545
Solidago altissima, 545
substrate-borne vibrations, 374
Mesoamerica,
biogeography, 1
new species, 1
Rutelidae, 1
Yaaxkumukia, 1
Metamasius callizona,
bromeliads, 1146
development, 1146
fecundity, 1146
weevil, 1146
Mexico,
bark beetle, 945
Dendroctonus, 945
genetic structure, 945
isozyme, 945
microbial control,
alate, 524
imported fire ant, 524
infection, 524
pathogen, 524
microsatellite,
colony genetic structure, 1101
eastern subterranean termite, 1101
Formosan subterranean termite, 1101
population genetics, 1101
microsporidia,
body size, 837
large aspen tortrix, 837
reproductive performance, 837
midgut, 592, 910, 917
bacteria, 917
cellularization, 592
Chrysoperla carnea, 917
cleavage, 592
embryogenesis, 592
energids, 592
larvae, 917
regeneration, 910
regenerative cells, 910
transmission electron miscroscopy, 910
yeast, 917
Zygentoma, 910

midwestern species,
instars, 755
nymphal key, 755
Sinea, 755
Miocene amber,
Brachypsectra, 409
fossil new species, 409
identification key, 409
relationships, 409
mitochondrial DNA, 157, 697, 959
Aedes vexans, 157
cytochrome oxidase subunit I, 959
genetic variation, 157, 697
phylogenetic analysis, 959
population structure, 157
Reticulitermes, 697
termite, 697
Tetranychus urticae, 959
mitochondrial DNA cytochrome oxidase I,
Aleyrodidae, 768
Asia, 768
Bemisia tabaci, 768
molecular marker, 768
moisture requirements,
Balaustium, 560
velvet mite, 560
water vapor uptake, 560
molecular diagnostics,
filth fly parasitoids, 799
Pteromalidae, 799
Spalangia, 799
molecular marker,
Aleyrodidae, 768
Asia, 768
Bemisia tabaci, 768
mitochondrial DNA cytochrome oxidase I, 768
molecular phylogenetics,
12S, 352
16S, 352
COII, 352
phylogenetics, 352
molecular systematics,
invasive pest, 195
Tsuga, 195
monitoring,
canola, 64
Ceutorhynchus obstrictus, 64
emergence, 64
sex ratio, 64
monophyly,
new species, 41
parasitoid, 41
phylogeny, 41
Siphonini, 41
morphology, 151, 433
cytochrome oxidase *c*, 433
Dioryctria, 433
DNA barcoding, 433
growth trajectory, 151
Lethocerus deyrolli, 151
mtDNA, 433
phenotypic plasticity, 151
predation, 151
morphometrics,
genitalic evolution, 144
intraspecific variation, 144
Phyllophaga hirticula, 144
postcopulatory sexual selection, 144
mtDNA, 433, 457
chromosome rearrangement, 457
cytochrome oxidase *c*, 433
Dioryctria, 433
DNA barcoding, 433
introgression, 457
morphology, 433
phylogeography, 457
Robertsonian fusion, 457
mutant,
eye color, 966
Hematobia irritans, 966
mutualism,
ants, 545
Membracidae, 545
Solidago altissima, 545
mycangial fungi,
bark beetle, 189
mycophagy, 189
symbiosis, 189
mycoheterotrophy,
Gastrodia, 851
orchid, 851
pollination, 851
Scatophagidae, 851
mycophagy,
bark beetle, 189
mycangial fungi, 189
symbiosis, 189
Myrmeciinae,
ant evolution, 487
Eocene, 487
Formicidae, 487
Okanagan Highlands, 487

N

Nemobiinae,
Allonemobius, 449
calling song, 449
Orthoptera, 449
phylogeny, 449
Neotropical Scarabaeidae,
biomass, 831
preservatives, 831
volume, 831
volume–mass relationship, 831
nest excavation,
Atta texana, 891
central cavity, 891
feeder holes, 891
fungus gardens, 891
Neuroptera,
Chrysopidae, 58
key, 58
Meleoma, 58
taxonomy, 58
new genus,
phylogeny, 1064
systematics, 1064
new species, 1, 41, 463
biogeography, 1, 463
diversity survey, 463
identification keys, 463
Mesoamerica, 1
monophyly, 41
parasitoid, 41
phylogeny, 41
Rutelidae, 1
Siphonini, 41
West Indies, 463
Yaaxkumukia, 1
New World,
cicada killer, 793
cicada-killer, 793
key, 793
Sphecius, 793
Nezara viridula,
enteric bacteria, 91
Enterococcus faecalis, 91
Klebsiella pneumoniae, 91
novel host,
buck moth, 1182
Hemileuca maia species complex, 1182
host shift, 1182
preference–performance, 1182
nutrient sink,
biocontrol, 845
Cecidochares, 845
Chromolaena, 845
gall, 845
nutrition,
ELISA, 321
food quality, 321
predation, 321
reproduction, 321
nymphal key,
instars, 755
midwestern species, 755
Sinea, 755
Nyssa,
Acronicta, 638
larval evolution, 638
wood-boring, 638

O

Okanagan Highlands,
ant evolution, 487
Eocene, 487
Formicidae, 487
Myrmeciinae, 487

oogenesis,
glassy-winged sharpshooter, 920
leafhopper, 920
sperm, 920
telotrophic, 920
Opiinae,
cultivated, 261
Fopius, 261
indigenous, 261
Psyttalia, 261
orchid,
Gastrodia, 851
mycoheterotrophy, 851
pollination, 851
Scatophagidae, 851
Oregon,
agricultural ecosystems, 33
crane fly pest species, 33
DNA diagnosis, 33
urban landscapes, 33
organogenesis,
Diatraea, 1206
embryonic stages, 1206
RNA levels, 1206
oriental fruit fly,
bulking agent, 1191
liquid diet rearing, 1191
sponge cloth, 1191
Oriental fruit moth,
autodetection, 898
female response, 898
Grapholita molesta, 898
sex pheromone, 898
Orius insidiosus,
prey defenses, 317
soybean, 317
soybean aphid, 317
Orthoptera,
Allonemobius, 449
calling song, 449
Nemobiinae, 449
phylogeny, 449
ovarian development,
color dimorphism, 1127
elytra, 1127
Lissorhoptrus oryzophilus, 1127
mating, 1127
oversized apparatus,
allometry, 327
body size, 327
intraspecific variability, 327
phylogenetic inertia, 327
overwintering,
bigeyed bug, 300
critical photoperiod, 300
sensitive stage, 300
oviposition,
eggs, 586
ELISA, 586
Parafilm, 586
predator, 586
ovipositional strategy,
behavior, 253
egg distribution, 253
stored-products, 253
Oxymonadida,
anaerobic protists identification, 1028
Parabasalia, 1028
symbiotic protists, 1028
termite identification, 1028

P

Pandemis pyrusana,
Choristoneura rosaceana, 617
courtship behavior, 617
Parabasalia,
anaerobic protists identification, 1028
Oxymonadida, 1028
symbiotic protists, 1028
termite identification, 1028
Parafilm,
eggs, 586
ELISA, 586
oviposition, 586
predator, 586
paralysis,
developmental delay, 1199
Melittobia digitata, 1199
venom, 1199
parasitism,
biological control, 113
Eulophidae, 113
high-fidelity polymerase chain reaction, 113
Platygastridae, 113
parasitoid, 41, 84, 1110
boreal forest, 1110
community composition, 1110
landscape structure, 1110
life table, 84
Malacosoma disstria, 1110
monophyly, 41
new species, 41
phylogeny, 41
reproduction, 84
Siphonini, 41
stored product pest, 84
pathogen,
alate, 524
imported fire ant, 524
infection, 524
microbial control, 524
PBAN,
calcium channel, 905
pheromone biosynthesis, 905
receptor signal transduction, 905
Persea americana,
Cales noacki, 553
demography, 553
fortuitous biological control, 553
jackknife, 553
Pheidole ceres,
ants, 398
food storage, 398
foraging, 398
seeds, 398
phenology,
Gonatocerus latipennis, 1154
leafhopper biology, 1154
Polynema sp., 1154
Vitis girdiana, 1154
phenotype,
asymmetry, 7
development, 7
larvae, 7
Leucochrysinae, 7
phenotypic plasticity,
growth trajectory, 151
Lethocerus deyrolli, 151
morphology, 151
predation, 151
pheromone biosynthesis,
calcium channel, 905
PBAN, 905
receptor signal transduction, 905
pheromones,
Prostephanus punctatus, 175
Prostephanus truncatus, 175
Rhyzopertha dominica, 175
Zelus tetracanthus, 175
pheromonostatic peptide,
gonad-specific virus, 969
Helicoverpa zea, 969
Hz-2V, 969
mating behavior, 969
phonoresponse,
acoustic behavior, 978
communication, 978
photoperiod,
Anthonomus grandis, 1085
boll weevil, 1085
reproductive diapause, 1085
Phyllophaga hirticula,
genitalic evolution, 144
intraspecific variation, 144
morphometrics, 144
postcopulatory sexual selection, 144
phylogenetic analysis,
cytochrome oxidase subunit I, 959
mitochondrial DNA, 959
Tetranychus urticae, 959
phylogenetic inertia,
allometry, 327
body size, 327
intraspecific variability, 327
oversized apparatus, 327

phylogenetics,
12S, 352
16S, 352
COII, 352
molecular phylogenetics, 352
phylogeny, 41, 241, 449, 723, 743, 1006, 1064
18s rDNA, 241
adult morphology, 723
Allonemobius, 449
Andex insignis, 743
androconia, 723
calling song, 449
cladistic analyses, 1006
Dytiscidae, 743
endophallus, 1006
Ephemeroptera, 241
Hyphydrini, 743
larvae, 743
monophyly, 41
Nemobiinae, 449
new genus, 1064
new species, 41
Orthoptera, 449
parasitoid, 41
Pristosia impunctata sp. nov., 1006
Pristosia latistoma sp. nov., 1006
Siphonini, 41
systematics, 1064
ultrasound, 723
phylogeography,
chromosome rearrangement, 457
introgression, 457
mtDNA, 457
Robertsonian fusion, 457
Phytoseiidae,
Anystidae, 536
biological control, 536
Erythraeidae, 536
Stigmaeidae, 536
Pierce's disease, 648, 879, 884
attachment, 884
biological control, 879
glassy-winged sharpshooter, 648
Gonatocerus tuberculifemur, 879
Graphocephala, 884
Homalodisca, 884
Homalodisca coagulata, 648, 879
Propetes, 648
Xylella fastidiosa, 648
pinene,
alkenal, 859
dorsal abdominal gland, 859
exuviae, 859
limonene, 859
pitcher plants,
Fletcherimyia, 218
Sarcophagidae, 218
Sarraceniaceae, 218
systematics, 218
plant morphology,
community ecology, 1133
detritus, 1133
indirect effect, 1133
tritrophic, 1133
plantains,
Cosmopolites sordidus, 875
head capsules width, 875
instars, 875
precipitation, 875
plasmodesmata,
cell wall, 305
host–parasite interface, 305
sink tissues, 305
transmission electron micrograph, 305
Platygastridae,
biological control, 113
Eulophidae, 113
high-fidelity polymerase chain reaction, 113
parasitism, 113
pollination,
Gastrodia, 851
mycoheterotrophy, 851
orchid, 851
Scatophagidae, 851
pollinator,
biological control, 96
Charles Robertson, 96
floral resources, 96
fly, 96
Polynema sp.,
Gonatocerus latipennis, 1154
leafhopper biology, 1154
phenology, 1154
Vitis girdiana, 1154
Ponerinae,
ant–host association, 567
Ectatomma, 567
Eucharitidae, 567
Kapala, 567
population genetics,
colony genetic structure, 1101
eastern subterranean termite, 1101
Formosan subterranean termite, 1101
microsatellite, 1101
population stability,
Cassida rubiginosa, 292
demographic traits, 292
mark–recapture, 292
reproductive rate, 292
population structure,
Aedes vexans, 157
genetic variation, 157
mitochondrial DNA, 157
postcopulatory sexual selection,
genitalic evolution, 144
intraspecific variation, 144
morphometrics, 144
Phyllophaga hirticula, 144
precipitation,
Cosmopolites sordidus, 875
head capsules width, 875
instars, 875
plantains, 875
predation, 151, 321
ELISA, 321
food quality, 321
growth trajectory, 151
Lethocerus deyrolli, 151
morphology, 151
nutrition, 321
phenotypic plasticity, 151
reproduction, 321
predator,
eggs, 586
ELISA, 586
oviposition, 586
Parafilm, 586
preference–performance,
buck moth, 1182
Hemileuca maia species complex, 1182
host shift, 1182
novel host, 1182
preservatives,
biomass, 831
Neotropical Scarabaeidae, 831
volume, 831
volume–mass relationship, 831
prey defenses,
Orius insidiosus, 317
soybean, 317
soybean aphid, 317
prezygotic isolation,
Canthon cyanellus, 1248
ecological speciation, 1248
sexual recognition, 1248
Prionus californicus,
mate location, 718
semiochemical, 718
wood borer, 718
prioritization,
impact, 121
introduced species, 121
invasive, 121
Pristosia impunctata sp. nov.,
cladistic analyses, 1006
endophallus, 1006
phylogeny, 1006
Pristosia latistoma sp. nov., 1006
Pristosia latistoma sp. nov.,
cladistic analyses, 1006
endophallus, 1006
phylogeny, 1006

Pristosia impunctata sp. nov., 1006
Propetes,
glassy-winged sharpshooter, 648
Homalodisca coagulata, 648
Pierce's disease, 648
Xylella fastidiosa, 648
Prostephanus punctatus,
pheromones, 175
Prostephanus truncatus, 175
Rhyzopertha dominica, 175
Zelus tetracanthus, 175
Prostephanus truncatus,
pheromones, 175
Prostephanus punctatus, 175
Rhyzopertha dominica, 175
Zelus tetracanthus, 175
Psyttalia,
cultivated, 261
Fopius, 261
indigenous, 261
Opiinae, 261
Pteromalidae,
filth fly parasitoids, 799
molecular diagnostics, 799
Spalangia, 799

Q

Qinling Mountains,
bamboo, 1057
eriophyids, 1057
key, 1057
taxonomy, 1057

R

rDNA,
5.8S, 369
Anastrepha suspensa, 369
ITS2, 369
Tephritidae, 369
receptor signal transduction,
calcium channel, 905
PBAN, 905
pheromone biosynthesis, 905
regeneration,
midgut, 910
regenerative cells, 910
transmission electron miscroscopy, 910
Zygentoma, 910
regenerative cells,
midgut, 910
regeneration, 910
transmission electron miscroscopy, 910
Zygentoma, 910
relationships,
Brachypsectra, 409
fossil new species, 409
identification key, 409
Miocene amber, 409
relative abundance index,
geographical distribution, 70
Mediterranean fruit fly, 70
Tephritidae, 70
remating,
dispersal, 607
mate choice, 607
sperm, 607
sterile insect technique, 607
Remington suture zone,
biogeography, 933
scanning electron microscopy, 933
specialization, 933
reproduction, 84, 321
ELISA, 321
food quality, 321
life table, 84
nutrition, 321
parasitoid, 84
predation, 321
stored product pest, 84
reproductive compatibility,
cryptic species complex, 387
sexual isolation, 387
sterile insect technique, 387
temporal isolation, 387
reproductive diapause,
Anthonomus grandis, 1085
boll weevil, 1085
photoperiod, 1085
reproductive performance,
body size, 837
large aspen tortrix, 837
microsporidia, 837
reproductive rate,
Cassida rubiginosa, 292
demographic traits, 292
mark–recapture, 292
population stability, 292
resistance management,
Bt corn, 279
insect dispersal, 279
southwestern corn borer, 279
transgenic crops, 279
Reticulitermes,
genetic variation, 697
mitochondrial DNA, 697
termite, 697
revision,
beetles, 776
Cybocephalus, 776
scale predator, 776
taxonomy, 776
RGB,
color analysis, 865
cotton, 865
image analysis, 865
vision, 865
Rhagoletis,
Diptera, 662
distribution, 662
host plant, 662
Tephritidae, 662
Rhyzopertha dominica,
pheromones, 175
Prostephanus punctatus, 175
Prostephanus truncatus, 175
Zelus tetracanthus, 175
RNA levels,
Diatraea, 1206
embryonic stages, 1206
organogenesis, 1206
Robertsonian fusion,
chromosome rearrangement, 457
introgression, 457
mtDNA, 457
phylogeography, 457
rough-leaved dogwood,
cicadas, 809
colonization, 809
heterogeneous habitats, 809
host plant density, 809
Rutelidae,
biogeography, 1
Mesoamerica, 1
new species, 1
Yaaxkumukia, 1

S

5.8S,
Anastrepha suspensa, 369
ITS2, 369
rDNA, 369
Tephritidae, 369
12S,
16S, 352
COII, 352
molecular phylogenetics, 352
phylogenetics, 352
16S,
12S, 352
COII, 352
molecular phylogenetics, 352
phylogenetics, 352
sandhills,
Attini, 673
Florida, 673
fungus gardening, 673
longleaf pine, 673
Sarcophagidae,
Fletcherimyia, 218
pitcher plants, 218
Sarraceniaceae, 218
systematics, 218
Sarraceniaceae,
Fletcherimyia, 218
pitcher plants, 218
Sarcophagidae, 218
systematics, 218
scale predator,
beetles, 776
Cybocephalus, 776

revision, 776
taxonomy, 776
scanning electron microscopy,
biogeography, 933
Remington suture zone, 933
specialization, 933
Scatophagidae,
Gastrodia, 851
mycoheterotrophy, 851
orchid, 851
pollination, 851
Scolytidae,
ambrosia beetles, 1034
exotic species, 1034
invasive pests, 1034
Xyleborini, 1034
Scolytinae,
body size, 211
Costa Rica, 211
host specificity, 211
Xyleborini, 211
secondary production,
Caenidae, 821
Ephemeroptera, 821
life history, 821
seeds,
ants, 398
food storage, 398
foraging, 398
Pheidole ceres, 398
semiochemical,
mate location, 718
Prionus californicus, 718
wood borer, 718
sensitive stage,
bigeyed bug, 300
critical photoperiod, 300
overwintering, 300
sex pheromone,
autodetection, 898
female response, 898
Grapholita molesta, 898
Oriental fruit moth, 898
sex ratio,
canola, 64
Ceutorhynchus obstrictus, 64
emergence, 64
monitoring, 64
sexual dimorphism,
alate, 625
developmental constraint, 625
eclosion, 625
swarming, 625
sexual isolation,
cryptic species complex, 387
reproductive compatibility, 387
sterile insect technique, 387
temporal isolation, 387
sexual recognition,
Canthon cyanellus, 1248
ecological speciation, 1248
prezygotic isolation, 1248
sexual selection, 342, 1244
allometry, 342
cricket, 1244
life history strategy, 1244
mate choice, 1244
Megasoma, 342
somatic proportions, 342
shore flies,
Hydrellia wirthi Korytkowski, 999
invasive species, 999
short-term alternative hosts,
Asphondylia, 1165
Cecidomyiidae, 1165
host range, 1165
life history strategy, 1165
Sinea,
instars, 755
midwestern species, 755
nymphal key, 755
sink tissues,
cell wall, 305
host-parasite interface, 305
plasmodesmata, 305
transmission electron micrograph, 305
Siphonini,
monophyly, 41
new species, 41
parasitoid, 41
phylogeny, 41
Solanaceae,
Ceratitis anonae, 530
Ceratitis fasciventris, 530
Ceratitis rosa, 530
Solenopsis invicta,
fire ants, 1213
genetic structure, 1213
invasion history, 1213
Solidago,
gall midges, 656
inquiline, 656
Solidago altissima,
ants, 545
Membracidae, 545
mutualism, 545
somatic proportions,
allometry, 342
Megasoma, 342
sexual selection, 342
southwestern corn borer,
Bt corn, 279
insect dispersal, 279
resistance management, 279
transgenic crops, 279
soybean,
Orius insidiosus, 317
prey defenses, 317
soybean aphid, 317
soybean aphid,
Orius insidiosus, 317
prey defenses, 317
soybean, 317
Spalangia,
filth fly parasitoids, 799
molecular diagnostics, 799
Pteromalidae, 799
specialization,
biogeography, 933
Remington suture zone, 933
scanning electron microscopy, 933
sperm, 607, 920
dispersal, 607
glassy-winged sharpshooter, 920
leafhopper, 920
mate choice, 607
oogenesis, 920
remating, 607
sterile insect technique, 607
telotrophic, 920
Sphecius,
cicada killer, 793
cicada-killer, 793
key, 793
New World, 793
sponge cloth,
bulking agent, 1191
liquid diet rearing, 1191
oriental fruit fly, 1191
stadium duration,
Gynaephora menyuanensis, 1012
head capsule width, 1012
instars, 1012
life cycle, 1012
stem borer,
biological weed control, 683
common reed, 683
life history, 683
wetlands, 683
sterile insect technique, 387, 607
cryptic species complex, 387
dispersal, 607
mate choice, 607
remating, 607
reproductive compatibility, 387
sexual isolation, 387
sperm, 607
temporal isolation, 387
Stigmaeidae,
Anystidae, 536
biological control, 536
Erythraeidae, 536
Phytoseiidae, 536
stinkbug,
Buchnera, 577
Heteroptera, 577
maximum likelihood, 577

symbiosis, 577
stored product pest,
life table, 84
parasitoid, 84
reproduction, 84
stored-products,
behavior, 253
egg distribution, 253
ovipositional strategy, 253
substrate-borne vibrations,
Ennya, 374
jamming, 374
Membracidae, 374
swarming,
alate, 625
developmental constraint, 625
eclosion, 625
sexual dimorphism, 625
symbiosis, 189, 577
bark beetle, 189
Buchnera, 577
Heteroptera, 577
maximum likelihood, 577
mycangial fungi, 189
mycophagy, 189
stinkbug, 577
symbiotic protists,
anaerobic protists identification, 1028
Oxymonadida, 1028
Parabasalia, 1028
termite identification, 1028
systematics, 218, 1064
Fletcherimyia, 218
new genus, 1064
phylogeny, 1064
pitcher plants, 218
Sarcophagidae, 218
Sarraceniaceae, 218

T

taxonomy, 58, 707, 776, 1057
Albitarsis Complex, 707
Anopheles marajoara, 707
bamboo, 1057
beetles, 776
Chrysopidae, 58
Cybocephalus, 776
distribution, 707
eriophyids, 1057
ITS2, 707
key, 58, 1057
Meleoma, 58
Neuroptera, 58
Qinling Mountains, 1057
revision, 776
scale predator, 776
telotrophic,
glassy-winged sharpshooter, 920
leafhopper, 920
oogenesis, 920
sperm, 920
temporal isolation,
cryptic species complex, 387
reproductive compatibility, 387
sexual isolation, 387
sterile insect technique, 387
Tephritidae, 70, 369, 662
5.8S, 369
Anastrepha suspensa, 369
Diptera, 662
distribution, 662
geographical distribution, 70
host plant, 662
ITS2, 369
Mediterranean fruit fly, 70
rDNA, 369
relative abundance index, 70
Rhagoletis, 662
termite,
genetic variation, 697
mitochondrial DNA, 697
Reticulitermes, 697
termite identification,
anaerobic protists identification, 1028
Oxymonadida, 1028
Parabasalia, 1028
symbiotic protists, 1028
Tetranychus urticae,
cytochrome oxidase subunit I, 959
mitochondrial DNA, 959
phylogenetic analysis, 959
trade-off,
Callosobruchus chinensis, 364
egg number, 364
egg size, 364
genetic correlation, 364
transgenic crops,
Bt corn, 279
insect dispersal, 279
resistance management, 279
southwestern corn borer, 279
transmission electron micrograph,
cell wall, 305
host–parasite interface, 305
plasmodesmata, 305
sink tissues, 305
transmission electron miscroscopy,
midgut, 910
regeneration, 910
regenerative cells, 910
Zygentoma, 910
tree hole,
abundance, 1172
bamboo, 1172
deforestation, 1172
diversity, 1172
Trialeurodes vaporariorum,
behavior, 598
Bemisia tabaci biotype A, 598
whitefly, 598
tritrophic,
community ecology, 1133
detritus, 1133
indirect effect, 1133
plant morphology, 1133
Tsuga,
invasive pest, 195
molecular systematics, 195

U

ultrasound,
adult morphology, 723
androconia, 723
phylogeny, 723
urban landscapes,
agricultural ecosystems, 33
crane fly pest species, 33
DNA diagnosis, 33
Oregon, 33

V

velvet mite,
Balaustium, 560
moisture requirements, 560
water vapor uptake, 560
venom,
developmental delay, 1199
Melittobia digitata, 1199
paralysis, 1199
vision,
color analysis, 865
cotton, 865
image analysis, 865
RGB, 865
Vitis girdiana,
Gonatocerus latipennis, 1154
leafhopper biology, 1154
phenology, 1154
Polynema sp., 1154
volume,
biomass, 831
Neotropical Scarabaeidae, 831
preservatives, 831
volume–mass relationship, 831
volume–mass relationship,
biomass, 831
Neotropical Scarabaeidae, 831
preservatives, 831
volume, 831

W

water vapor uptake,
Balaustium, 560
moisture requirements, 560
velvet mite, 560
weevil,
bromeliads, 1146
development, 1146

fecundity, 1146
Metamasius callizona, 1146
West Indies, 164, 463
biogeography, 463
citrus pest, 164
diversity survey, 463
genetic markers, 164
Hispaniola, 164
identification keys, 463
invasive species, 164
new species, 463
wetlands,
biological weed control, 683
common reed, 683
life history, 683
stem borer, 683
whitefly,
behavior, 598
Bemisia tabaci biotype A, 598
Trialeurodes vaporariorum, 598
wood borer,
mate location, 718
Prionus californicus, 718
semiochemical, 718
wood-boring,
Acronicta, 638
larval evolution, 638
Nyssa, 638

X

Xyleborini, 211, 1034
ambrosia beetles, 1034
body size, 211
Costa Rica, 211
exotic species, 1034
host specificity, 211
invasive pests, 1034
Scolytidae, 1034
Scolytinae, 211
Xylella fastidiosa,
glassy-winged sharpshooter, 648
Homalodisca coagulata, 648
Pierce's disease, 648
Propetes, 648

Y

Yaaxkumukia,
biogeography, 1
Mesoamerica, 1
new species, 1
Rutelidae, 1
yeast,
bacteria, 917
Chrysoperla carnea, 917
larvae, 917
midgut, 917

Z

Zelus tetracanthus,
pheromones, 175
Prostephanus punctatus, 175
Prostephanus truncatus, 175
Rhyzopertha dominica, 175
Zygentoma,
midgut, 910
regeneration, 910
regenerative cells, 910
transmission electron miscroscopy, 910